

Drug Facts

Active ingredient (in each 0.5 ml) Purpose

Racinephrine, USP 11.25mg.....Bronchodilator
(as 13.5 mg of Racinephrine HCl, USP).

Uses

For temporary relief of mild symptoms of intermittent asthma:

- Wheezing • Tightness of chest • Shortness of breath

Warnings

Asthma alert: Because asthma may be life threatening, see a doctor if you:

- Are not better in 20 minutes • Get worse
- Need more than 12 inhalations in 24 hours
- Use more than 9 inhalations in 24 hours for 3 or more days a week
- Have more than 2 asthma attacks in a week

These may be signs that your asthma is getting worse.

Do not use:

- Unless a doctor said you have asthma
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.
- If product is brown in color or cloudy
- If its color is pinkish or darker than slightly yellow or if it contains a precipitate.

Ask a doctor before use if you have:

- Ever been hospitalized for asthma • Heart disease
- High blood pressure • Diabetes • Thyroid disease
- Seizures • Narrow angle glaucoma
- A psychiatric or emotional condition
- Trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are:

- Taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions.
- Taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain).

When using this product:

- **Your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death.**
- **Your risk of heart attack or stroke increases if you:**
 - **Have a history of high blood pressure or heart disease**
 - **Take this product more frequently or take more than the recommended dose.**
- Avoid foods or beverages that contain caffeine
- Avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect. →

Drug Facts (continued)

Warnings (continued)

Stop use and ask a doctor if:

- Your asthma is getting worse (See Asthma alert)
- You have difficulty sleeping
- You have a rapid heartbeat
- You have tremors, nervousness, or seizure
- **If pregnant or breast-feeding**, ask a health professional before use.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **Do not use more than directed**
- **For use in a hand-held rubber bulb nebulizer:**
Add 0.5 ml (contents of one vial) of solution to nebulizer.
- Adults and children 4 years of age and over:
1 to 3 inhalations not more often than every 3 hours.
Do not use more than 12 inhalations in 24 hours.
The use of this product by children should be supervised by an adult.
- Children under 4 years of age: ask a doctor.

Other Information

- Protect from light. Avoid excessive heat. Protect from freezing.
- Store between 2°C and 25°C (36°F and 77°F).
- How Supplied: S₂[®] Racinephrine Inhalation Solution, USP 2.25% is supplied in 0.5 ml sterile unit-of-use vials.
NDC 0487-5901-99 30 vials, each vial in an individual foil pouch.

Inactive ingredients

Ascorbic acid, citric acid, edetate disodium, water for injection.

Questions? 1-800-443-4313

Monday through Friday, 8:00 a.m. to 5:00 p.m.
(Eastern Standard Time).